


FROM

TUTTLE, MOREHOUSE & TAYLOR,
Printers and Bookbinders,
New Haven, Ct.



Q.L. 567.1 Glover, Townsend
~~Set~~ Illustrations of North American
Glover Entomology
Washington, 1872

MBL/WHOI



0 0301 0047622 2

Revised

Picture & Text.

ILLUSTRATIONS
OF
NORTH AMERICAN ENTOMOLOGY,
(UNITED STATES AND CANADA,)

BY TOWNEND GLOVER, WASHINGTON, D. C.

ORTHOPTERA.

CONTENTS:

Introduction.....	III
Arrangement of Families.....	V
Plates.....	
Notes on Food and Habits of Orthoptera.....	1
Parasites.....	2
List of Substances Injured by Orthoptera.....	5
List of Genera figured.....	7
List of Species figured.....	9
List of Desiderata and Errata..	"

*With Compliments of Townsend Glover.
(Please acknowledge receipt.)*

Entered according to Act of Congress, in the year 1872, by
TOWNEND GLOVER,
in the Office of the Librarian of Congress at Washington, D. C.

INTRODUCTION.

It is not the design of the author in the following work, to present scientific or highly-finished engravings of North American Orthoptera, but merely figures, giving a general idea of their form, size, and color, to aid the young Entomologist in the identification of species. At the same time reference is made to their names in MR. SAMUEL H. SCUDDER's "Catalogue of Orthoptera of North America," published by the Smithsonian Institution, Washington, 1868, and to his "Materials for a Monograph of the North American Orthoptera," published in "Proceedings Boston Society Natural History, 1862," where many original descriptions as well as the names of their authors may be found. The REV. CYRUS THOMAS has described many new species in the "Proceedings of the Academy of Natural Sciences," Philadelphia, 1870, and in the "Geological Surveys of Territories, made by DR. F. V. HAYDEN," 1870 and 1871.

In some cases it has been found necessary to figure European insects, where no specimens of the allied genera or species could be procured in the United States. In all such instances, the figures will be marked as European, and due credit given to the original work from which they were taken.

In the arrangement of the list accompanying each plate, the first name given is always that of the latest and best authority; and the last, the name of the genus in which it was formerly placed by its first discoverer. As the works of SAY and HARRIS are most widely disseminated in the Northern and Eastern states, their names will also be given as synonyms when deemed necessary.

Some of the insects figured have been transferred from other orders to the Orthoptera, for the reasons given below, namely:—The *Thripidae* were formerly placed by WESTWOOD by themselves in *Thysanoptera*, and more lately by DR. PACKARD among the *Hemiptera*: but as WESTWOOD distinctly says that "as they possess two setiform *mandibles*, and the disposition of the other parts of the mouth are those of a *real mandibulated insect*," they cannot properly be retained amongst the haustellate Hemiptera, and are therefore placed provisionally amongst the Orthoptera. *Podura* and its allies also possess small *mandibles* with minute teeth, and are not mentioned by MR. SCUDDER in his list of Orthoptera; but as LEUNIS in his "Synopsis

der drei Naturreiche" places them under the head of "Division B, false or wingless Orthoptera," they have also been classed with the Orthoptera until a better place can be found for them. The *Mallophaga* or bird lice also possess *distinct jaws* instead of a sucking tube, (Verrill,) and have small and hook-like mandibles. Such being the case, they cannot be placed with the true *Pediculina* which are decidedly haustellate, and are, therefore, also placed provisionally in Division B, or Wingless Orthoptera of LEUNIS.

The text to accompany these plates will consist of merely the latest name, and some of the principal synonyms of the insects figured, with a concise history of their habits as far as known; their food, habitat; the best remedies used to destroy them if injurious, and the name of the donor, with a very short description of the principal families, or genera, and in some instances a brief notice of the principal difference, or any striking peculiarity, which distinguishes closely allied species from each other, so as to call attention to that particularly in the comparison or arrangement of specimens in a cabinet. There will also be a full alphabetical list of the principal vegetable and animal substances injured by any species at the end of the volume, for the use of farmers who do not understand anything about Entomology, but who can thus be enabled to identify any particular insect by referring to the plant on which it is generally found feeding.

The thanks of the author are due to MR. S. I. SMITH, of Yale College, New Haven, Connecticut; MR. CHAS. R. DODGE, of Washington, D. C.; MR. C. V. RILEY, of St. Louis, Mo. for specimens kindly given from their own private collections, and from which the original drawings were prepared. Especial acknowledgment is due to the REV. CYRUS THOMAS, of Illinois, and MR. PHILIP R. UHLER, of Baltimore, Md., not only for type specimens, but also for their valuable aid in comparing them with others, and in determining doubtful species. It may also be well to observe, that the original type specimens have been carefully preserved in the Entomological cabinet of the Museum of the Department of Agriculture in Washington, for future reference by any Entomologist who may wish to work up or study the Orthopterous insects of the United States and Canada.

This small pioneer edition of fifty copies is intended for distribution to Entomologists and Entomological Societies *only*. The plates have been etched and the text written either after or before the hours of official duty; and it is also published at the expense of the author, and not by the Department of Agriculture. If it should be approved of, it is proposed to publish yearly, or from time to time additional plates &c. of the same size and in similar style of any new or rare Orthoptera which may be added to our list by the exploring expeditions, or by private enterprise, as likewise eventually to illustrate all the other orders of insects in a similar manner.

INTRODUCTION.

V

The following table of the principal families of Orthoptera has been temporarily arranged for the use of young Entomologists who wish to classify their collections. The name of the family to which each insect belongs in SCUDDER'S Catalogue, &c. will be placed (in italics) as the end of the list of names and synonyms appended to each plate.


Division A. True Orthoptera.	Gryllides. Scudder.	{	Ex. <i>Acanthus</i> , or flower cricket. <i>Gryllus</i> , com-		
	<i>Achetidae</i> (<i>Leach</i>) of <i>Westwood</i> .		mon cricket, and <i>Gryllotalpa</i> or mole cricket.		
	Locustariae. Scudd.	{	Phaneroptera, Phylloptera, Microcentrum and Cyr-		
	<i>Gryllidae</i> . (<i>Leach</i> .) <i>Westw.</i>		tophyllus. (Katydid.) <i>Xyphidium</i> ; <i>Orcheli-</i> mum, &c.; slender meadow grasshoppers or Katydid, &c.; (<i>Ceuthophilus</i> .) large wingless cricket or grasshopper of Utah, and <i>Hadenæ-</i> cus, cave wingless cricket.		
	Acrydii. Scudd.	{	<i>Aceridium</i> , <i>Caloptenus</i> , <i>Edipoda</i> , <i>Stenobothrus</i> , &c.;		
	<i>Locustidae</i> . (<i>Leach</i> .) <i>Westw.</i>		true grasshoppers; and <i>Tettix</i> , grouse locusts.		
Division B. False Orthoptera	Phasmida. Scudd.	{	Spectres. Walking sticks.		
	<i>Phasmidae</i> . <i>Westw.</i>				
	Mantides. Scudd.	{	Rear horses or soothsayers.		
	<i>Mantidae</i> . <i>Westw.</i>				
	Blattariae. Scudd.	{	Cockroaches.		
	<i>Blattidae</i> . (<i>Stephens</i> .) <i>Westw.</i>				
Forficulariae. Scudd.	{	Earwigs.			
<i>Forficulidae</i> . (<i>Stephens</i> .) <i>Westw.</i>					
Division B. False Orthoptera	Fringed wings.	{	Thripidae	{	formerly order <i>Thysanoptera</i> of <i>West-</i> <i>wood</i> .
	Poduridae	{	Spring and bristle tails.	{	formerly order <i>Thysanura</i> , <i>Burm.</i> , and lately in <i>Neuro-</i> <i>ptera</i> . <i>Packard</i> .
	Wingless. Orthoptera	{	Mallophaga.	{	in order <i>Orthoptera</i> by <i>Gerstaecker</i> , &c., but placed in <i>Hemiptera</i> by <i>Packard</i> .

The False Orthoptera have been put after the True Orthoptera, so that they may either be left out altogether, or removed into other orders if found desirable.

PLATE I.---ORTHOPTERA.

Fig. 1	{ Platamodes (Scudd.) pennsylvanica. Scudd. } Blatta (Linn.) " De Geer. }	*Scudd. Cat. 70	Scudd. Mono. 117	<i>Blattariae</i>
2	{ Periplaneta (Burm.) americana. Burm. } Blatta (Linn.) " Linn. }	62.	"	116
3	{ Platamodes (Scudd.) pennsylvanica. Scudd. } Blatta (Linn.) " De Geer. }	70	"	117
4	{ Ectobia (Westw.) germanica. Stephens. } Blatta (Linn.) " Aude & Brulle. }	11	"	117
5	{ Stylopyga (Fischer) orientalis. Fisch. deW. } Blatta (Linn.) " Linn. }	78	"	116
6	" " " ♀ "	78.	"	116.
7	{ Diapheromera (Gray) femorata. Scudd. } Bacteria (Latr.) Sayii. Burm. } Bacunculus (Burm.) femoratus. Uhler. } Spectrum (Serv.) femoratum. Say. }	11.	"	0. <i>Phasmaria</i>
8	{ Anisomorpha (Scudd.) buprestoides. Gray. } Spectrum (Serv.) bivittatum. Say. } Phasma (Licht.) buprestoides. Stoll. }	9.	"	0.
9	Stenopelmatus (Burm.) talpa. ♂ Burm.	78.	"	0. <i>Locusturiae</i>
10	Labia (Leach) minuta. Scudd.	13.	"	115. <i>Forficulariae</i>
11	Thamnotrizon (Fischer) dorsale. ♀ Burm.	0.	"	0. <i>Locusturiae</i>
12	Gryllotalpa (Latr.) longipennis. ♂ Scudd.	29.	"	126. <i>Gryllides</i>
13	{ Gryllus (Linn.) pennsylvanicus. ♀ Burm. } Acheta (Fab.) pennsylvanica. Uhler. }	36.	"	128. "
14	" " " ♂ "	36.	"	128. "
15	{ Acridium (Geoff.) americanum. ♀ Scudd. } Cyrtacanthacris (Walk.) " Walk. } Gryllus (Linn.) " Drury. }	1.	"	166. <i>Acrydia</i>
16	{ Caloptenus (Serv.) bivittatus. ♀ Uhler. } " " femoratus. Burm. } Gryllus (Linn.) bivittatus. Say. }	20.	"	165. "

*Scudder's Catalogue and Monograph, see Introduction.



J. Glover

PLATE II.---ORTHOPTERA.

MANTIS CAROLINA (Linn.) Scudder's Catalogue, p. 48. *Phasmida*

- Fig. 1 Egg case.
 2 Egg case with young escaping.
 3 Young Mantis devouring another.
 4 Young Mantis in the Nymph state (Gray variety.)
 5 Young " " " " (Green variety.)
 6 Imago. Mantis carolina. (Linn.) ♂
 7 " " " " ♀ (Gray var.)
 8 " " " " ♀ (Green var.)
 9 Egg case cut longitudinally to exhibit structure.
 10 " " " " "
 11 " " transversely " "
 12 Wings of ♂
 13, 14, 15 Base of Antennae, fore legs and tarsi.
 16 Wings of ♀
 * 17 Sciara?

* This fly was popularly believed to be connected with the yellow fever which raged in Norfolk, Va. in 1855. They were exceedingly numerous there during the whole time the yellow fever was prevalent in that year, and hence were known in Norfolk and its vicinity as "yellow fever flies." (See Sciara in Diptera.)



J. Sliver

PLATE III.---ORTHOPTERA.

Fig. 1	{	Edipoda (Latr.) eucrata.	Uhler. }	Scudd. Cat	56.	Scudd. Mono.	472.	<i>Acrigdu</i>
	{	Locusta (Linn.)	" Harr. }					
2	"	"	" ♀ "	"	56.	"	472.	"
3	{	Edipoda (Latr.) discoidea.	♂ Serv. }	"	56.	"	469.	"
	{	Acridium (Geoff.) tuberculatum.	Pal de }					
	{	Beauv.						
4	{	Romalea (Serv.) microptera.	+ Serv. }	"	71.	"	0.	"
	{	Gryllus (Fab.) centurio.	Burm. }					
5	{	Ceuthophilus (Scudd.) maculatus.	♀ Scudd. }	"	21.		434.	<i>Locustinae.</i>
	{	Rhaphidophora (Charp.) maculata.	Harr. }					
	{	Phalangopsis (Burm.)	" Harr. }					
	{	Ephippigera (Serv.)	" Say. }					
	{	MSS.						
6		Tridactylus (Oliv.) terminalis.	Scudd.	"	82.	"	425.	<i>Gryllides.</i>
7	{	Edipoda (Latr.) discoidea.	♀ Serv. }	"	56.	"	468.	<i>Acrigdu.</i>
	{	Acridium (Geoff.) tuberculatum.	Pal de }					
	{	Beauv.						
8		Edipoda (Latr.) corallipes.	♀ Hald	"	56.	"	0.	"
9	{	Nemobius (Serv.) vittatus.	♀ Scudd. }	"	54.	"	430.	<i>Gryllides.</i>
	{	Acheta (Fab.) servilis.	Harr. and Say MSS. }					
10	"	"	♂ " "	"	54.	"	430.	"
11		Orocharis (Uhler.) saltator.	♀ Uhler.	"	58.	"	0.	"
12	"	"	♂ "	"	58.	"	0.	"




PLATE IV.---ORTHOPTERA.

Fig. 1	{ Ecantus (Serv.) niveus. ♀ Dr. Geer. " " fasciatus. Fitch. Gryllus (Linn.) niveus. Oliv. }	Scudd. Cat. 55. Scudd. Mono. 431. <i>Gryllides.</i>
2	" " " ♂ "	" 55. " 431.
3	{ Microcentrum (Scudd.) retinervis. ♀ Scudd. Phylloptera (Serv.) " Burm. }	" 53. " 436. <i>Locustariæ.</i>
4	{ Phylloptera (Serv.) oblongifolia ♀ Burm. Locusta (Linn.) " De Geer. }	" 68. " 445. "
5	{ Ecantus (Serv.) bipunctatus. ♀ De Geer. Ecantus " punctulatus. Fitch. Gryllus (Linn.) bipunctatus. De Geer. }	" 55. " 432. <i>Gryllides.</i>
6	" " " ♀ "	" 55. " 432. "
7	{ Orchelimum (Serv.) vulgare. ♀ Harr. Pterophylla (Kirby MSS.) agilis. Harr. }	" 59. " 452. <i>Locustariæ.</i>
8	" " " ♂ "	" 59. " 452. "
9	{ Mesops (Serv.) Wyomingensis. Thomas. Opomala (Serv. emend.) Wyomingensis. Thomas. }	Proceedings Academy Natural Science. Philadelphia, 1871. <i>Acridia.</i>
10	{ Nyphidium (Serv.) fasciatum. ♂ Serv. Pterophylla (Kirby MSS.) fasciata. Harr. Orchelimum (Serv.) gracile. Harr. Locusta (Linn.) fasciata. De Geer. }	Scudd. Cat. 84 Scudd. Mono. 451. <i>Locustariæ.</i>
11	Nyphidium (Serv.) brevipennis. ♀ Scudd.	" 84. " 451. "
12	{ Conocephalus (Thunb.) ensiger. ♀ Harr. Locusta (Linn.) acuminata. Stoll. }	" 22. " 449. "
13	Stenobothrus (Fisch.) admirabilis. Uhler.	" 77. " 0. <i>Acridia.</i>
14	{ Pyrgomorpha (Fisch.) brevicornis. ♀ Thomas. Opomala (Serv. emend.) " " Truxalis (Linn.) " Fab. Gryllus (Linn.) (Acridium Fab.) brevi- cornis. Linn. }	" 83. " 0. "
15	{ Cyrtophyllus (Burm.) concavus. ♂ Scudd. Platyphyllum (Serv.) concavum. Harr. Pterophylla (Kirby MSS.) concava. Say and Harr. }	" 24. " 444. <i>Locustariæ.</i>
16	{ Egg and young of Microcentrum retin- ervis. Scudd. See Fig. 3. }	" 53. " 436. "



J. Glover.

PLATE V.---ORTHOPTERA.

Fig. 1	*Tettix (Latr. emend.) ornata. Scudd.				Scudd. Cat. 79. Scudd. Mono. 474. <i>Aerydii</i> .
	Tetrix (Latr.) quadrimaculata. ♀ Harr.				
	Var. figured.				
	Tetrix (Latr.) bilineatus. Harr. Var.				
	" " dorsalis. " "				
2	" " sordida. " "				" 79. " 474. "
	" " arenosa. Burm. "				
	Acridium (Geoff.) ornatum. Say				
	" " " " "				
	" " " " "				
3	{ Edipoda (Latr.) carolina. Burm. }				" 56. " 468. "
	{ Locusta (Linn.) caroliniana. Catesby. }				
	{ Locusta (Linn.) corohna. Linn. & Harr. }				
4	{ Edipoda (Latr.) phoenicoptera. ♀ }				" 57. " 468. "
	{ Germ. Locusta (Linn.) corallina. Harr. }				
5	{ Acridium (Geoff.) rubiginosum. ♀ Harr. }				" 7. " 467. "
	{ MSS. Acridium (Geoff.) danificum. Sauss. }				
6	{ Edipoda (Latr.) sulphurea. ♂ Burm. }				" 47. " 470. "
	{ Gryllus (Fab.) " Fab. }				
	{ Locusta (Linn.) " Linn & Harr. }				
7	{ Tettix (Latr. emend.) lateralis. Scudd. }				" 79. " 477. "
	{ Tetrix (Latr.) " Harr. }				
	{ Acrydium (Geoff.) " Say. }				
8	{ Batrachidea (Serv.) cristata. Scudd. (Var.) }				" 11. " 478. "
	{ Tetrix (Latr.) " Harr. MSS. }				
9	{ Tragocephala (Harr.) viridilasciata. ♀ }				" 82. " 461. "
	{ Harr. }				
	{ Tragocephala (Harr.) radiata. Harr. }				
	{ Edipoda (Latr.) virginiana. Burm. }				
10	{ Acridium (Geoff.) virginianum. Oliv. }				" 65. " 448. <i>Locustarue</i>
	{ Egg of Phaneroptera (Serv.) curvicauda. Serv. }				
	{ Phaneroptera (Serv.) angustifolia. Harr. }				
11	{ Locusta (Linn.) curvicauda. De Geer. }				" 20. " 464. <i>Aerydii</i> .
	{ Caloptenus (Serv.) femur rubrum. ♀ }				
12	{ Burm. }				" 6. " 467. "
	{ Acridium (Geoff.) femur rubrum. Harr. }				
13	{ Acridium (Geoff.) obscurum. ♀ Burm. }				" 56. " 0. "
	{ Gryllus (Fab.) }				
14	{ Edipoda (Latr.) eucerata. Uhler in }				" 77. " 458. "
	{ Harr. Florida variety. }				
	{ Locusta (Linn.) Harr. }				
15	{ Stenobothrus (Fischer.) maculipennis. ♀ }				" 0. " 457. "
	{ Scudd. }				
16	{ Stenobothrus (Fischer.) longipennis. ♂ }				" 20. " 465. "
	{ Scudd. }				
	{ Caloptenus (Serv.) bivittatus. ♂ Uhler. }				
17	{ " " femoratus. Burm. }				" 20. " 465. "
	{ Gryllus (Linn.) bivittatus. Say. }				

*All these so-called species of Harris, &c., are said (by Scudder) to be merely varieties of *Tettix* ornata, (Scudd.) (*Acridium* ornatum, Say.) and that the *T. ornata* of Harris is the *T. granulatus* of Scudder's monograph, p. 414.



J. Flower

PLATE VI.---ORTHOPTERA.

Fig. 1	Philopterus (Nitzsch.) falciicornis. Nitzsch. Europe. From *Lennis, p. 650. Pediculus (Linn.) payonius. Linn. Eu- rope. From Lennis.							Mallophaga.
2	Machilis (Latr.) variabilis. Say. Massachusetts.							Apteros. } Orthoptera. }
3	Machilis (Latr.) variabilis. Say. Massachusetts.							"
4	Phloeothrips (Halid.) coriacea. Halid. From Westwood.							Thripida.
5	Thrips (Linn.) sp.? Europe. From Lennis, p. 618.							"
6	Lepisma (Linn.) saccharina. Linn. From Lennis, p. 650.							Apteros. } Orthoptera. }
7	Podura (Linn.) villosa. Fab. Europe. From Lennis, p. 650.							"
	7a. Smynthurus (Latr.) sp.? Not described (Maryland, in fungi)							"
8	Trichodectes (Nitzsch.) latus. De Geer. Europe. From Lennis, p. 651							Mallophaga.
9	Thrips (Linn.) sp.? (Maryland, in greenhouse.)							Thripida
10	{ Chloeahtis (Harr.) conspersa. ♂ Harr. } Chloeahtis (Harr.) abortiva. Harr. }	Scudd. Cat. 22	Scudd. Mono. 455.					Acryphi
11	{ Chloeahtis (Harr.) viridis. ♀ Scudd. } Stenobothrus (Fischer.) viridis. ♀ Scudd. }	" 22.	" "				155.	"
12	Phylloptera (Serv.) rotundifolia. ♀ Scudd.	" 69.	" "				115.	Locustar.
13	{ Nemobius (Serv.) fasciatus. ♀ Scudd. } Acheta (Fab.) hospes. Fab. } Gryllus (Fab.) fasciatus. De Geer. }	" 55.	" "				130.	Gryllides.
14	{ Gryllus (Fab.) domesticus. ♀ Pack. } Acheta (Fab.) domestica. Thomas. Ill. }	" 2.	" "				0.	Acryphi
15	{ Stenobothrus (Fischer.) curtippennis. ♂ Scudd. } Chloeahtis (Harr.) curtippennis. Harr. } Locusta (Linn.) " " }	" 77.	" "				456.	"
16	Pezotettix (Burm.) borealis. ♀ Scudd.	" 63.	" "				464.	"
17	" " " ♂ " "	" 63.	" "				461.	"
18	" " " ♀ " "	" 63.	" "				464.	"
19	Forficula (Linn.) sp.? From N. Y.							Forficular.
20	Cryptocercus (Scudd.) punctulatus. Scudd.	" 24.	" "				420.	Blattaria.
21	Stenobothrus (Fisch.) aequalis. ♀ Scudd.	" 77.	" "				459.	Acryphi.
22	Phyllopalpus (Uhler.) pulchellus. Uhler.	" 68.	" "				0.	Gryllides.
23	Oedipoda (Latr.) euerata. ♂ Erichs.	" 56.	" "				472.	Acryphi.
24	{ Opomala (Serv. emend.) bivittata. ♀ Serv. } Opsomala (Serv.) bivittata. Serv. } Acridium (Geoff.) " De Haan. }	" 58.	" "				0	"
25	Phrynotettix verruculata (Uhler. MSS.) From Pecos River, Texas.	"	"				0	"
26	{ Opomala (Serv. emend.) bivittata ♂ Serv. } Opsomala (Serv.) " " }	" 58.	" "				0	"
27	Stenobothrus (Fischer.) maculipennis. ♀ var. Scudd.	" 76.	" "				0	"

*Lennis. "Synopsis der drei Naturreiche." Hannover.




PLATE VII.---ORTHOPTERA.

Fig. 1	{ Tettigidea (Scudd.) polymorpha. Scudd. Tetrix (Latr.) parvopennis. Harr. " " polymorpha. Burm (Var. B.) }	Scudd. Cat. 79 Scudd. Mono. 477. <i>Acrydii</i> .
2	{ Phaneroptera (Serv.) curvicauda. ♂ (Serv.) Phaneroptera (Serv.) angustifolia. Harr. Locusta (Linn.) curvicauda. De Geer. }	" 65. " 448. <i>Locustariac.</i>
3	" " " ♀ "	" 65. " 448. "
4	{ Ceuthophilus (Scudd.) lapidicolus ♀ (Scudd.) Rhaphidiphora (Charp.) lapidicola. Burm. Phalangopsis (Serv.) " Burm. }	" 21. " 435. "
5	" " " "	" 21. " 435. "
6	{ Orchelimum (Serv.) vulgare. ♂ Harr. Pterophylla (Kirby MSS.) agilis. Harr. }	" 57. " 452. <i>Gryllides</i> .
7	" " " ♀ "	" 57. " 452. "
8	{ Copiophora (Walk.) mucronata ♂ Thomas. (Hayden's Geol. Surv. Terr. 1871; 444, and Can. Ent. 1872. Copiophora (Serv. of Agass.) }	" 23. " 0. <i>Acrydii</i> .
9	{ Edipoda (Latr.) marmorata. ♀ Uhler. Locusta (Linn.) " Harr. }	" 56. " 473. "
10	{ Stenobothrus (Fischer.) curtipennis. ♂ Scudd. Chloealtis (Harr.) curtipennis ♂ Harr. Locusta (Linn.) " Harr. }	" 77. " 456. "
11	{ Brachyepplus (Charp.) magnus. ♀ Girard. }	" 20. " 0. "
12	{ Stylopyga (Fisch.) orientalis. ♀ Fischer, Blatta (Linn.) }	" 78. " 416. <i>Blattariac.</i>
13	{ Gryllotalpa (Latr.) borealis. ♂ Burm. " " brevipennis. Serv. " " americana. Say. MSS. " " borealis. Burm. }	" 29. " 456. <i>Gryllides</i> .
14	Daibina (Hald.) brevipes. ♂ Hald.	" 24. " 443. <i>Locustariac.</i>
15	" " " ♀ "	" 24. " 443. "
16	{ Anabrus (Hald.) haldemanni. ♀ Girard. Pterolepis " Thomas. }	" 3. " 0. "
17	{ Gryllus (Linn.) abbreviatus. ♀ Serv. Acheta (Fab.) abbreviata. Harr. }	" 30. " 427. <i>Gryllides</i> .
18	{ Nemobius (Serv.) exiguus. ♀ Scudd. Acheta (Fab.) exigua. Say. }	" 55. " 429. "

VII





PLATE VIII.---ORTHOPTERA.

Fig. 1 Caloptenus (Serv.) spretus. ♀ Uhler MSS. Scudd. Cat. O. Scudd. Mono. O. Acrydia.					
2	{ Caloptenus (Serv.) femur-rubrum. ♀ Burm. Acridium (Geoff.) " " Harr. }	"	20.	"	464. "
3	{ Edipoda (Latr.) atrox. { Scudd. in Hay- den's Geol. Surv. Neb. 253. Hayden's Geol. Surv., 1871, 458. }				"
4	{ Pezotettix (Burm.) picta. ♀ Thomas. Proc. Acad. Nat. Scien., Phil. 1870. }				"
5	{ Thamnotrizou (Fischer.) trilineatus. ♀ Thomas. Hayden's Geol. Surv. Terr., 1871, p. 443. Decticus (Serv.) trilineatus. }				Locustariae.
6	{ Hadenæus (Scudd.) subterraneus. ♀ Scudd. Rhaphidophora (Charp.) subterraneus. }	"	40.	"	440. "
7	{ Tettix (Latr. emend.) lateralis(?) ♀ Scudd. Tettix (Latr.) " Harr. Acridium (Geoff.) " Say. }	"	79.	"	477. Acrydia.
8	{ Ceuthophilus (Scudd.) Uhleri. ♀ (affinis.) Scudd. }	"	21.	"	435. Locustariae.
9	{ Udeopsylla (Scudd.) robusta. ♂ Scudd. Phalangopsis (Serv.) " Hald. Daihinia (Hald.) " }	"	83.	"	443. "
10	{ Boopidon (Thomas.) flavofasciatum. ♀ Thomas. Proc. Acad. Nat. Sci. Phil., 1870. }				Acrydia.
11	{ Stauronotus (Fischer.) Elliotti. ♀ Thomas. Fr. Acad. Nat. Scien. Phil. 1870, 82. }				"
12	{ Caloptenus (Serv.) differentialis. ♀ Thomas. }	"	5.	"	0. "
13	{ Acridium (Geoff.) alutaceum. ♀ Harr. (Small spec.) Acridium rusticum (Burm. only.) " torvum. Say. and Harr. }	"	4.	"	466. "
11	{ Copiphora (Walk.) mucronata. ♀ Thomas. Hayden's Geol. Surv. Terr., 1871, p. 444. Copiphora (Serv.) }	"	23.	"	0. "
15	Camptonotus (Uhler.) Scudder. ♀ Uhler.	"	21.	"	Locustariae.

VIII



J. Glover

PLATE IX.---ORTHOPTERA.

Fig. 1	{ Anabrus (Hald.) simplex. ♀ Hald. Thomas. Hayden's Geol. Surv. Terr. 1871, p. 438. }	Scudd. Cat	8.	Scudd. Mono	0.	<i>Locustarius</i>
2	{ Oedipoda (Latr.) tenebrosa. ♀ Scudd. Thomas Geol. Surv. Terr. 1871, p. 459. Tomonotus (Sauss.) mexicanus. Thomas. Pr. Acad. Nat. Sci., Phil., 1870. }				0.	<i>Acrydia</i> .
3	{ Brachypeplus (Charp.) virescens. ♀ Charp. }	"	20.	"	0.	"
4	{ Caloptenus (Serv.) differentialis ♂ Thomas. }	"	5.	"	0.	"
5	Gryllus (Linn.) formosus. ♀ Say.	"	34.	"	0.	<i>Gryllodes</i> .
6	{ Oedipoda (Latr.) tritasciata. ♀ Walk. Thomas. Hayden's Geol. Surv. Terr. 1871, p. 456. Oedipoda (Latr.) pruinosa. Thomas. Pr. Acad. Nat. Sci., Phil., 1870. Gryllus (Linn.) trifasciata. Say. Am. Ent. Fig. 3, pl. 34. }	"	39.	"	0.	<i>Acrydia</i> .
7	{ Acrolophitus (Thomas.) lurtipes ♀ Thomas. Gryllus (Linn.) birtipes. Say. }	"	38.	"	0.	"
8	{ Decticus (Serv.) pallidipalpis. ♀ Thomas. Hayden's Geol. Surv. Terr. 1871, p. 442. }	"	24	"	0.	<i>Locustarius</i>
9	{ Locusta (Linn.) fuliginosa. ♂ Thomas. Hayden's Geol. Surv. Terr. 1871, p. 443. }					<i>Acrydia</i> .
10	{ Gryllus (Linn.) luctuosus. ♀ Barm. and Serv. Acheta (Linn.) luctuosus. }	"	35.	"	427	"
11	Acheta (Linn.) luctuosus ♂	"	35.	"	427.	"

IX



J. Glover

PLATE X.--ORTHOPTERA.

Fig 1	{ Diapheromera (Gray) femorata Scudd. Bacteria (Latr.) sayii Burm. Baeneculus (Burm.) femoratus Uhler. Spectrum (Serv.) femoratum, Say. }	Scudd. Cat. 11	Scudd. Mono.	Phasmida
2	{ Porficesila (Latr.) { From Westwood. } gigantea Serv. { European. }	"	27.	" 0. <i>Porficesilariae</i>
3	{ Labia (Leach) minor. Doubled. } Porficsila (Linn.) minor. Burm. }	"	43.	" "
4	{ Stenobothrus (Fischer) curtipennis. ♂ Scudd. Chloealtis (Harr.) curtipennis. Harr. Locusta " " Scudd. }	"	77.	" 456. <i>Aerypha</i>
5	{ Chloealtis (Harr.) viridis. ♀ Scudd. Stenobothrus (Fisch.) viridis }	"	22.	" 445 "
6	{ Edipoda (Latr.) verruculata. ♂ Scudd. Locusta (Linn.) latipennis. Harr. " " verruculata Kirby. }	"	57.	" 471. "
7	{ Edipoda (Latr.) carinata. ♀ Scudd. Tr. Am. Ent. Soc., Vol. 2, p. 306. }	"		" "
8	Gryllus (Linn.) neglectus. ♂ ♀ Scudd.	"	36.	" 428. <i>Gryllodes</i>
9	{ Phylloptera (Serv.) oblongifolia. ♂ Burm. Locusta (Linn.) oblongifolia. De Geer. }	"	68.	" 445. <i>Locustariae</i>
10	{ Tragocephala (Harr.) infuscata ♂ Harr. Gomphocerus (Thunb.) infuscata. Uhler. }	"	82.	" 461. <i>Aerydia</i>
11	{ Edipoda (Latr.) sordida. ♂ Burm. Locusta (Linn.) nebulosa. Harr. }	"	57.	" 473. "
12	{ Chloealtis (Harr.) conspersa. ♀ Harr. " " abortiva. " }	"	22	" 455. "
13	Aeridium (Geoff.) alutacenm. ♀ Harr.	"	4	" 466. "
14	{ Stenobothrus (Fisch.) maculipennis ♀ Scudd. }	"	77	" 458 "





PLATE XI.---ORTHOPTERA.

Fig. 1 *Acridium* (Geoff.) *frontalis*. ♀ Thomas. Hayden's Geol. Surv. Terr., 1871, p. 448.

- | | | |
|----|---|---------------------|
| 2 | $\left\{ \begin{array}{l} \text{Caloptenus (Serv.) occidentalis. } \text{♀} \\ \text{Thomas. Hayden's Geol. Surv. Terr.,} \\ \text{1871, p. 453.} \end{array} \right\}$ | <i>Acrydi.</i> |
| 3 | $\text{Caloptenus (Serv.) viridis. } \text{♀} \text{ Thomas. Hayden's Geol. Surv. Terr., 1871, p. 450.}$ | " |
| 4 | $\left\{ \begin{array}{l} \text{Caloptenus (Serv.) Dodgei. } \text{♂} \text{ Thomas.} \\ \text{Hayden's Geol. Surv. Terr., 1871,} \\ \text{p. 451.} \end{array} \right\}$ | " |
| 5 | $\left\{ \begin{array}{l} \text{Caloptenus (Serv.) Dodgei. } \text{♂} \text{ Thomas.} \\ \text{Hayden's Geol. Surv. Terr., 1871,} \\ \text{p. 451.} \end{array} \right\}$ | " |
| 6 | $\left\{ \begin{array}{l} \text{Caloptenus (Serv.) differentialis. } \text{♂} \\ \text{Thomas. (Dark var.)} \end{array} \right\}$ | Scudd. Cat. 5. |
| 7 | $\left\{ \begin{array}{l} \text{Opomala (Serv. emend.) bivittata. } \text{♀} \\ \text{Say.} \\ \text{Opomala (Serv.)} \end{array} \right\}$ | " 58. |
| 8 | $\left\{ \begin{array}{l} \text{Mesops (Serv.) Wyomingensis. } \text{♀} \\ \text{Thomas. Proc. Phil. Acad. Nat. Sci.,} \\ \text{1871.} \\ \text{Opomala (Serv. emend.)} \end{array} \right\}$ | " |
| 9 | $\text{Caloptenus (Serv.) Dodgei. } \text{♀} \text{ Thomas. (See above.)}$ | " |
| 10 | $\left\{ \begin{array}{l} \text{Caloptenus (Serv.) Turnbullii. } \text{♂} \text{ Thomas.} \\ \text{(Var. A.) Hayden's Geol. Surv. Terr.,} \\ \text{1871, p. 452.} \end{array} \right\}$ | " |
| 11 | $\left\{ \begin{array}{l} \text{Ephippitytha (Serv.) gracilipes. } \text{♂} \\ \text{Thomas.} \end{array} \right\}$ | <i>Locustariae.</i> |
| 12 | $\text{Phaneroptera (Serv.) coloradensis. } \text{♂} \text{ Thomas. MSS.}$ | " |
| 13 | $\left\{ \begin{array}{l} \text{Pezotettix (Burm.) obesa. } \text{♀} \text{ Thomas.} \\ \text{Hayden's Geol. Surv., Terr., 1871,} \\ \text{p. 454.} \end{array} \right\}$ | <i>Acrydi.</i> |
| 14 | $\text{Pezotettix (Burm.) obesa. } \text{♀} \text{ Thomas.}$ | " |
| 15 | $\text{Ephippigera (Serv.) tschivavensis. } \text{♀} \text{ (Hald.) Stansbury's Report.}$ | " |
| 16 | $\left\{ \begin{array}{l} \text{Locusta (Linn.) occidentalis. } \text{♀} \text{ Thomas.} \\ \text{Hayden's Geol. Surv. Terr., 1871, p.} \\ \text{444.} \end{array} \right\}$ | " |
| 17 | $\left\{ \begin{array}{l} \text{Pterolepis (Serv.) minutus. } \text{♀} \text{ Thomas.} \\ \text{Hayden's Geol. Surv. Terr., 1871, p. 441.} \\ \text{Anabrus (Hald.) minutus. Thomas.} \end{array} \right\}$ | <i>Locustariae.</i> |

XI




T. Flevers

PLATE XII.---ORTHOPTERA.

- Fig. 1 Nirmus argulus. From Rev. J. G. Wood. Nat. Hist., illustrated, p. 686. *Mallophaga.*
- 2 Menapon pallidum. Nat. Hist., illustrated, p. 686. "
- 3 Trichodectes longicornis. Nat. Hist., illustrated, p. 685. "
- 4 Trichodectes equi. Nat. Hist. illustrated, p. 685. "
- 5 Docophorus cygni. Nat. Hist., illustrated, p. 686. "
- 6 Haematopinus piliferus. Nat. Hist., illustrated, p. 685. "
- 7 { Haematopinus suis. Denny Monog. an-
opulurum. Nat. Hist., illustrated, p. }
685. }
- 8 Edipoda (Latr.) rugosa. ♀ Scudd. Scudd. Cat. 55. Scudd. Mono. 469. *Acrydii.*
- 9 Thrips (Linn.) cerealeum. Halid. From an English work. *Thripidae.*
- 10 " " " " " " " "
- 11 Phleothrips autumnalis. Uhler. Found on Oak, Md. "
- 12 { Edipoda (Latr.) montana. ♀ Thomas. }
Hayden's Geol. Surv. Terr., 1871, p. 462. } *Acrydii.*
- 13 { Edipoda (Latr.) cincta. ♀ Thomas. }
Proc. Acad. Nat. Sci., Phil., 1870. 70. }
Hayden's Geol. Surv. Terr., 1871, p. 461. } "
- 14 { Caloptenus (Serv.) griseus. ♀ Thomas. }
Hayden's Geol. Surv. Terr., 1871, p. 454. } "
- 15 { Edipoda (Latr.) undulata. Thomas. }
Hayden's Geol. Surv. Terr., 1871, p. 460. } "
- 16 Mantis (Linn.) New sp. *Mantides.*
- 17 { Edipoda (Latr.) maritima. ♂ Scudd. }
Locusta (Linn.) maritima. (Harr.) } Scudd. Cat. 56. Scudd. Mono. 472. *Acrydii.*
- 18 { Stenobothrus (Fischer.) curtipennis. ♂ }
Scudd. } " 77. " 456. "
Chlœaltis (Harr.) curtipennis. Harr. }
Locusta (Linn.) " Harr. }
- 19 { Tettix (Latr. emend.) ornata. Scudd. }
Acridium (Fab.) ornatum. Say. (From }
Say's fig.) }
20. Edipoda (Latr.) pellucida. ♀ Scudd. " 57. " 472. "
- 21 { Edipoda (Latr.) carlingiana. ♂ Thomas. }
Proc. Acad. Nat. Sci., Phil., 1870, 81, and }
Hayden's Geol. Surv. Terr., 1870, p. 275. }

XII



J. Glover

PLATE XIII.---ORTHOPTERA.

- Fig. 1 { Eggs of Phylloptera (Serv.) oblongifolia. }
 Burm. } Scudd. Cat. 68. Scudd. Mono. 445. *Locustariac.*
 Locusta (Linn.) oblongifolia. De Geer. }
- 2 Pezotettix (Burm.) nebrascensis. Thomas. Hayden's Geol. Surv. Terr., 1871, p. 455. *Acrydii.*
- 3 { Edipoda (Latr.) Haldemanni. Scudd. Hayden's Geol. Surv., Neb., p. 253. " }
 { Edipoda (Latr.) corallipes. Hald. Stansbury's Rep., p. 371, pl. 10, fig. 2. }
- 4 Edipoda (Latr.) longipennis. Thomas. Hayden's Geol. Surv., 1871, p. 463. "
- 5 Batrachidea (prob.) pupa. From Md. "
- 6 { Thamnotrizon (Fischer) scabricollis. Thomas. Hayden's Geol. Surv. Terr., *Locustariac.*
 1871, p. 441. }
- 7 { Stenopelmatus (Burm.) fasciatus. Thomas. (large sp. from Cal.) Hayden's Geol. " }
 Surv. Terr., 1871, p. 434. }
- 8 { Edipoda (Latr.) aequalis. Uhler. }
 Locusta (Linn.) " Harr. } Scudd. Cat. 55. Scudd. Mono. 470. *Acrydii.*
 Gryllus (Linn.) " Say. }
- 9 Pezotettix (Burm.) unicolor. Thomas. "
- 10 { Boopidon (Thomas) nubilum. Thomas. Hayden's Geol. Surv. Terr., 1870, p. 273. " }
 Boopidon (Thomas) nigrum. Thomas. Gryllus (Linn.) nubilus. Say. }
- 11 Mantis (Linn.) missouriensis. Riley. MSS. *Mantides.*
- 12 Mantis (?) Thomas. From Wyoming Terr. "
- 13 Oxycoryphus (Fischer) obscurus. Thomas. Hayden's Geol. Surv. Terr., 1871, p. 466. *Acrydii.*
- 14 { *Acridium (Geoff.) semi-rubrum. Saussure. Scudd. Cat. 7. " }
 Acridium (Geoff.) flavicorne of Stoll. }
- 15 { Caloptenus (Serv.) spretus. Uhler. (Young insect.) Thomas. " }
 Desc. Hayden's Geol. Surv. 1871, p. 451. }

*Acridium semi-rubrum is said, by Rev. Cyrus Thomas, to have been taken in Southern Texas, but is mentioned by Fitch in Trans. N. Y. S. Ag. Soc., 1856, p. 490, as occurring in Cayenne only.

XIII



J. H. V. 1860

XIV



J. Flower



F. Steyer.

NOTES ON FOOD
AND
HABITS OF ORTHOPTERA.

The list of vegetable and animal substances injured by Orthoptera, as given below, comprises a very small number of the most important, as it would take up too much space in a merely preliminary sketch to enumerate them all.

Should, however, the text relating to the insects figured be published, it is designed to give a full list of all the substances injured, as well as all the insects injuring them, as mentioned by the various authors who have written on the subject.

Before commencing the list it may, however, be advisable to mention some of the principal families, in relation to their food, in general terms. The *Gryllina* (*Ceuthophilus*, Pl. III, Fig. 5; *Ceanthus*, Pl. IV, Figs. 1, 2,) are mostly vegetable feeders, whilst *Anabrus simplex* (Pl. IX, Fig. 1,) varies its diet, according to the REV. CYRUS THOMAS, by devouring the Cicada, or harvest-flies, whenever it can capture them.

The true crickets (*Gryllus* or *Acheta*, Pl. X, Fig. 8,) live under stones, &c., and, although vegetable feeders, living on roots and herbage, yet, in many instances, they injure animal substances, such as cloth, leather, &c., when it is in the neighborhood of their haunts. The Mole crickets (*Gryllotalpa*, Pl. VII, Fig. 13,) burrow under the earth like moles, and usually feed on roots and herbage, and in Europe do considerable injury. The Locustariae (*Phaneroptera*, Pl. VII, Figs. 2, 3; *Cyrtophyllus*, Pl. IV, Fig. 15, katydids,) inhabit shrubs and trees, feeding principally on their foliage, whilst the slender meadow grasshopper, *Orchelimum* (Pl. IV, 78,) feeds mostly on grass and herbage. The wingless Locustariae (*Stenopelmatus*, Pl. XIII, Fig. 7,) live on the ground and are said to feed on vegetable substances.

The Acrydii, or true grasshoppers and locusts, (*Caloptenus*, Pl. VII, *Acridium*, Pl. I, Fig. 15; *Edipoda*, Pl. V, Fig. 3; and *Locusta*, Pl. IX, Fig. 9,) live principally amongst the grass and low herbage, and devour all kinds of vegetable substances. Some species assemble in great numbers when migrating, like the locust of scripture, and are extremely injurious to almost all kinds of vegetation. The Mantides (*Mantis carolina*, Pl. II,) are altogether predaceous. They kill and devour all other insects they can over-

power, and are, therefore, almost the only family amongst the Orthoptera which are at all beneficial to the farmer as destroying other injurious insects. The Phasmidae, spectres, or walking sticks, (*Diapheromera*, Pl. X, Fig. 1,) resemble twigs, in form and color, and feed on the foliage of various shrubs and trees, whilst the Blattariae, or cockroaches, (*Stylopyga*, Pl. VII, Fig. 12; *Ectobia*, Pl. I, Figs. 4, 5,) are almost omnivorous, feeding indiscriminately on both vegetable and animal substances, and are extremely destructive in houses and on shipboard. The Forficulariae, or earwigs, (*Forficula*, Pl. X, Fig. 2,) are not numerous enough in this country to do much injury, but in Europe are said to injure flowers and ripe fruit, and occasionally to feed on small insects. The Thripidae certainly do much injury to the tender foliage of the grape vine, &c., and are very destructive to greenhouse plants. Thrips cerealeum, (Pl. XII, Figs. 9, 10,) injures the wheat crop in Europe by sucking the sap of the grain or seed and cause it to shrivel up and wither. MR. WALSH, however, considered the true Thrips in the light of a beneficial insect, stating that it destroyed the larvae of the destructive wheat midge *Cecidomyia* (*Diplosis*) *tritici* and other injurious insects.

Some of the wingless orthoptera (*Podura*, the springtail, Pl. VI, Fig. 17,) are stated, by MR. CURTIS, to feed upon the pulp of leaves, at the same time poisoning the sap and thus injuring the plants. The Mallophaga (or bird lice, Pl. XII, Fig. 1,) *Nirmus argulus* and dog louse, *Trichodectes latus*, Pl. VI, Fig. 8,) feed upon the feathers and hairs of the birds or animals they infest, and do not injure them by sucking the blood. By thus studying the food of the various Orthopterous insects, we find that all of them are more or less injurious to the farmer, with the sole exception of the Mantis, or rear-horse, as it is commonly called in Maryland and Virginia, which lives altogether on animal food, and is, therefore, beneficial as destroying injurious insects.

The family of Orthoptera is stated, by DR. PACKARD, to contain "about 5,000 species" and constitutes a very important article of food for certain animals, birds and fishes. Some of the grasshoppers are even eaten by our western Indians, when partially roasted and coarsely pulverized between two stones. A specimen of food used by the Indians now in the Museum of the Department of Agriculture, in Washington, sent from Camp Harney, Oregon, by Asst. Surgeon MOFFATT, U. S. A., which contains a great number of the heads and femora of some great wingless grasshopper or cricket, (apparently *Anabrus simplex*, Pl. IX, Fig. 1,) which is so abundant in some parts of the west. Several of the Orthoptera are also subject to be destroyed by both external and internal parasites. A coleopterous insect, *Rhipiphorus*, (*Symbius*) *blattarum*, probably the same as *Rhipidius*, mentioned by PACKARD, is parasitic in the bodies of certain cockroaches. Several species of Ichneumon-flies, *Evania loevigata*, Oliv, and a species of *Pteromalus* are said to destroy the eggs of cockroaches. A large species of *Asilus*, or robber-fly, according to the REV. C. THOMAS, also destroys numbers of grasshoppers by sucking out their juices. A species of *Tachina*, a two-winged fly, very similar

to our house fly, but larger, and having a more hairy body, destroys the perfect insect of the Mantis, or rear-horse, as many as nine having come out of the body of one *Mantis carolina*, (Pl. II, Figs. 1, 2, 3.) An Ichneumon-fly destroys the eggs of katydids. Crickets, grasshoppers and katydids are infested with the *Filaria*, *Gregarina* and *Gordins*, hair-snakes, or worms, which live in their bodies. A small scarlet-red mite: *Astoma locustarum* (WALSH) or *Ocypete* of HARRIS is frequently found clustered on the body or under the wings of grasshoppers, and it is said that when numerous, they eventually kill the insect they infest. In late summer and autumn great numbers of dead and dried-up grasshoppers are frequently observed in Maryland and Virginia, clinging fast to the tops of the highest stalks of grass or weeds. These probably have been destroyed by some animal or vegetable parasite, at present unknown.

Many of the so-called dirt-daubers, mud and sand wasps, provision their nests with young grasshoppers to serve as food for their larva, and there is no doubt that when more attention is given to the habits of our Orthoptera, many other parasites especially among the Ichneumon-flies, Chalcididea, &c., will be discovered and made known to the public.

ALPHABETICAL LIST

OF

VEGETABLE AND ANIMAL SUBSTANCES

INJURED BY

ORTHOPTERA.

The number of the plate on which the insect is figured will be distinguished by being in Roman numerals, whilst the number of the figure will be placed in italics, thus IV, 5; plate 4, figure 6.

When the mark, †, is placed before the name of an insect, it signifies that it is injurious, ††, very injurious; ‡, beneficial; †‡, more injurious than beneficial; and ‡†, more beneficial than injurious.

- Aphides. †‡*Ecanthus niveus*, IV, 1, 2, said to destroy plant lice, and also injures grapevines, &c.
- Beneficial. See Predaceous.
- Birds. *Nirmus argulus*, XII, 1. Louse. Parasitic upon
- Blackberry. †*Ecanthus niveus*, IV, 1, 2, deposits eggs in cane, and injures
- Cherry. †*Ecanthus niveus*, IV, 1, 2, deposits eggs in cane, and injures
- Books. †*Ectobia germanica*, I, 4, eats, destroys and covers with filth.
- Books. †*Stylopyga orientalis*, VII, 2, (and other cockroaches in general,) eats, destroys, and covers with filth.
- Caves. *Hadenococcus subterraneus*, VIII, 6, found in
- Clothing. †*Ectobia germanica*, I, 4, eats and destroys
- Clothing. †*Stylopyga orientalis*, VII, 2, eats and destroys
- Corn. See Maize.
- Currant. †*Ecanthus niveus*, IV, 1, 2, deposits eggs on twigs, and injures
- Deer fallow. *Trichodectes longicornis*, XII, 3 Louse, parasite on
- Diplosis tritici. See wheat midge.
- Dogs. *Trichodectes latus*, VI, 8. Louse, parasite on
- Dogs. *Haematopinus piliferus*, XII, 6. parasite on
- Fruits. †*Gryllus neglectus*, (and crickets in general,) X, 3, injure
- Fruits. †*Caloptenus femur-rubrum*, (and grasshoppers in general,) VIII, 2, injure
- Fungi. *Smynthurus*, VI, 7, inhabits
- Gnaphalium. *Anisomorpha buprestoides*, I, 5, probably feeds on
- Grain. ††*Caloptenus femur-rubrum* (and grasshoppers in general,) VIII, 2, destroys plants
- Grain. ††*Thrips cerealeum* (Europe.) XII, 2, 10, sucks out milky juice, and destroys
- Grape. †*Ecanthus niveus*, IV, 1, 2, devours foliage, severs branches, and injures shoots by depositing eggs in them.
- Grape. †*Orocharis saltator*, III, 11, 12, deposits eggs on branches, and injures
- Grass. ††*Caloptenus femur-rubrum*, (and all the grasshoppers in general,) VIII, 2, eat and destroy
- Grass. †*Edipoda*, sulphurea, &c., &c., V, 6, eat and destroy
- Grass. †*Acridium americanum*, &c., I, 15, (and many of the other orthoptera.)
- Greenhouse plants. ††*Thrips*. (?) VI, 9, injure
- Herbage. See insects destroying grass.
- Hog. *Haematopinus suis*, XII, 7, louse parasitic on
- Horse. *Trichodectes equi*, XII, 4, louse parasitic on
- Insects in general. See predaceous.
- Leather. †*Ectobia germanica*, (and other cockroaches,) I, 4, eat holes in
- Lemon. †*Phylloptera oblongifolia*, IV, 4, (and other insects found on orange,) injure foliage.
- Maize. ††*Caloptenus femur-rubrum*, VIII, 2, (and many of the other grasshoppers,) injure young plants and blades. See grass.
- Melons. †*Gryllus neglectus*, X, 3, (and other crickets,) injure

- Oak. *Acridium rubiginosum*, V, 8, found on
- Oak. *Phloethrips autumnalis*, XII, 11, found on
- Onion. †† *Limothrips tritici* (for Thrips see VI, 9,) injures
- Orange. † *Phylloptera oblongifolia*, IV, 4, injures foliage.
- Orange. † *Romalea microptera*, III, 4, injures foliage.
- Orange. † *Acridium obscurum*, V, 8, injures foliage.
- Palmetto. *Anisomorpha buprestoides*, I, 8, found on
- Paper. See books.
- Peacock. *Philopterus falcicornis*, VI, 1, louse parasitic on
- Peach. † *Ecanthus niveus*, IV, 1, 2, injures twigs by depositing eggs in them.
- Plum. † *Ecanthus niveus*, IV, 1, 2, injures twigs by depositing eggs in them.
- Potato. † *Gryllus neglectus*, X, 8, (and other crickets,) feeds on, and injures
- Potato. † *Caloptenus femur rubrum*, VIII, 2, (and other grasshoppers,) sometimes injure foliage.
- Potato. † *Gryllotalpa longipennis*, I, 12, (and mole crickets in general,) injure roots, &c.
- Poultry. *Menapon pallidum*, XII, 2, louse, parasitic on.
- Predaceous. ‖ *Mantis carolina*, II, 1, 2, feed entirely on other insects.
- Predaceous. †‖ *Forficula*, VI, 19, said to eat aphides, &c., but also destructive to flowers, fruits, &c., in Europe.
- Predaceous. †‖ Thrips, VI, 9, said to destroy larvæ of wheat midge, &c.
- Predaceous. †‖ *Anabrus simplex*, IX, 1, devours harvest-fly or cicada.
- Raspberry. † *Ecanthus niveus*, IV, 1, 2, injures twigs by depositing eggs in them.
- Roots in general, † *Gryllus neglectus*, X, 8, (and crickets in general,) feed upon and injure
- Roots in general. † *Gryllotalpa longipennis*, I, 12, (and mole crickets in general,) feed upon and injure
- Silk. † *Lepisma saccharina*, VI, 6, said to eat holes in
- Shrubs. † *Cyrtophyllus concavus*, IV, 12, (and katydids in general,) feed on
- Shrubs. *Diapheromera femorata*, X, 1, feeds on foliage.
- Squash. † *Gryllus neglectus*, X, 8, (and other crickets,) feeds upon, and injures
- Pumpkin, &c. † *Caloptenus femur rubrum*, VIII, 2, (and other grasshoppers,) injures foliage.
- Sugar-cane. *Gryllotalpa didactyla* in West Indies, injures
- Swan. *Dacophorus cygni*, louse parasitic on
- Trees. See shrubs.
- Vegetables. † *Gryllus neglectus*, (X, 8, (and other crickets,) injures roots and leaves.
- Vegetables. † *Gryllotalpa longipennis*, I, 12, (and other mole crickets,) injures roots and leaves.
- Vegetables. †† *Caloptenus femur rubrum*, VIII, 2, (and other grasshoppers,) destroys
- Vegetable substances. *Podura*, VI, 7, feeds upon
- Victuals. †† *Ectobia germanica*, I, 4, (and all cockroaches,) injures and render filthy.
- Wheat. †† *Thrips cerealum*, IX, 10, injures grain.
- Wheat midge. †‖ Thrips, V, 9, said by WALSH to destroy larvæ of wheat midge so destructive to grain
- Woollen cloth. † *Gryllus domestica*, VI, 14, eats holes in

ALPHABETICAL LIST OF GENERA FIGURED.

NOTE.—The number of the plate on which the insect is figured will be distinguished by being in Roman numerals, whilst the number of the figure will be in italics, thus: IV, 6; plate 4, figure 6.

- Acheta. See Gryllus, Nemobius.
 Acridium alutaceum, VIII, 13; X, 13.
 Acridium americanum, I, 15.
 Acridium frontalis, XI, 1.
 Acridium obscurum, V, 2.
 Acridium rubiginosum, V, 5.
 Acridium semirubrum, XIII, 14.
 Acridium. See also. Caloptenus, Edipoda, Opomala, Tettix, Tragocephala.
 Acrolophitus hirtipes, IX, 7.
 Anabrus. See also Pterolepis.
 Anabrus haldemannii, VII, 17.
 Anabrus simplex, IX, 1.
 Anisomorpha buprestoides, I, 3.
 Bacteria. See Diapheromera.
 Bacunculus. See Diapheromera.
 Batrachidea (pupa) XIII, 5.
 Batrachidea crestatata, V, 8.
 Blatta. See Ectobia Periplaneta.
 Platamodes, Stylopyga.
 Boopidon flavofasceatum, VIII, 10.
 Boopidon nubilum, XIII, 10.
 Brachypeplus magnus, VII, 11.
 Brachypeplus virescens, IX, 3.
 Caloptenus bivittatus, I, 15; V, 16.
 Caloptenus differentialis, VIII, 2; IX, 4; XI, 6.
 Caloptenus dodgei, XI, 4, 5, 9.
 Caloptenus femur-rubrum, V, 11; VIII, 2.
 Caloptenus griseus, XII, 14.
 Caloptenus occidentalis, XI, 2.
 Caloptenus spretus, VIII, 1; XIII, 15.
 Caloptenus turnbullii, XI, 10.
 Caloptenus viridis, XI, 3.
 Camptonotus scudderi, VIII, 15.
 Ceuthophilus lapidiculus, VII, 4, 5.
 Ceuthophilus maculatus, III, 5.
 Ceuthophilus Uhlerii, VIII, 8.
 Chlealtis conspersa, VI, 11; X, 12.
 Chlealtis viridis, X, 5.
 Chlealtis. See also Stenobothrus.
 Conocephalus ensiger, IV, 12.
 Copiophora mucronata, VII, 5; VIII, 14.
 Copiophora. See Copiophora.
 Cryptocercus punctulatus, VI, 20.
 Cyrtacanthacris. See Acridium.
 Cyrtophyllus concavus, IV, 15.
 Daibinia brevipes, VII, 14, 15.
 Daibinia. See Udeopsylla.
 Decticus. See also Thamnotrizon.
 Decticus pallidipalpis, IX, 3.
 Diapheromera femorata, I, 7; X, 1.
 Docophorus cygni, XII, 5.
 Ectobia germanica, I, 4.
 Ephippigera tschivavensis, XI, 15.
 Ephippigera. See also Ceuthophilus.
 Ephippitytha gracilipes, XI, 11.
 Forficesila gigantea, X, 2.
 Forficula. See also Labia.
 Forficula. (?) VI, 19.
 Gomphocerus. See Tragocephala.
 Gryllotalpa borealis, VII, 13.
 Gryllotalpa longipennis, I, 12.
 Gryllus abbreviatus, VII, 17.
 Gryllus domesticus, VI, 14.
 Gryllus (?) formosus, IX, 5.
 Gryllus luctuosus, IX, 10.
 Gryllus neglectus, X, 8.
 Gryllus pennsylvanicus, I, 13, 14.
 Gryllus. See also Acridium, Acrolophitus, Caloptenus, Ceanthus, Edipoda, Pyrgomorpha, Romalea.
 Hadenæus subterraneus, VIII, 6.
 Haematopinus piliferus, XII, 6.
 Haematopinus suis, XII, 4.
 Labia, minor, X, 3.
 Labia minuta, I, 10.
 Lepisma saccharina, VI, 6.
 Locusta fuliginosa, IX, 9.
 Locusta occidentalis, XI, 15.
 Locusta. See also Conocephalus, Gryllus, Edipoda, Phaneroptera, Phylloptera, Stenobothrus, Niphidium.
 Machilis variabilis, VI, 2, 3.
 Mantis (?) XII, 16.
 Mantis (?) XIII, 13.
 Mantis carolina, II, 1, 2, 3, &c.
 Mantis missouriensis, XIII, 11.
 Menopon pallidum, XII, 2.
 Mesops Wyomingsensis, IV, 9; XI, 8.
 Microcentrum retinervis, IV, 3.
 Microcentrum egg, IV, 16.
 Nemobius exiguius, VII, 18.
 Nemobius fasciatus, III, 9, 10.
 Nemobius vittatus, VI, 13.
 Nirmus argulus, XII, 1.
 Ceanthus bipunctatus, IV, 5, 6.

- Cecanthus niveus*, IV, 2.
Edipoda aequalis, XIII, 3.
 " *atrox*, VIII, 3.
 " *carinata*, X, 7.
 " *carolina*, V, 3.
 " *carlingiana*, XII, 21.
 " *cincta*, XII, 13.
 " *coralipes*, III, 8.
 " *discoidea*, III, 3, 7.
 " *eucrata*, III, 2; V, 13; VI, 25.
 " *Haldemanni*, XIII, 3.
 " *longipennis*, XIII, 4.
 " *maratima*, XII, 17.
 " *marmorata*, VII, 2.
 " *montana*, XII, 12.
 " *pellucida*, XII, 20.
 " *phœnicoptera*, V, 4.
 " *rugosa*, XII, 8.
 " *sordida*, X, 11.
 " *sulphurea*, V, 5.
 " *tenebrosa*, IX, 2.
 " *trifasciata*, IX, 3.
 " *verruculata*, X, 5.
 " *undulata*, XII, 15.
 " See also *Tragocephala*.
Opomala bivittata, VI, 24 and 26; XI, 7.
Opsomala. See *Opomala*, *Mesops* and *Pyrgomorpha*.
Orchelimum vulgare, IV, 7, 8; VII, 6.
Orchelimum. See also *Xiphidium*.
Orocharis saltator, III, 11, 12.
Oxycoryphus obscurus, XIII, 13.
Pediculus. See *Phyllopterus*.
Periplaneta americana, I, 2.
Pezotettix borealis, VI, 16.
 " *nebrascensis*, XIII, 2.
 " *obesa*, XI, 13, 14.
 " *picta*, VIII, 4.
 " *unicolor*, XIII, 9.
Phalangopsis. See *Ceuthophilus*, *Udeopsylla*.
Phaneroptera coloradensis, XI, 12.
 " *curvicauda*, VII, 2, 3, and V, 10.
Phyllopterus falcicornis, VI, 1.
Phlœthrips coreacea, VI, 4.
Phrynotettix verruculata, VI, 25.
Phylloptera (?) egg, IV, 16.
Phylloptera oblongifolia, X, 9; IV, 4; XIII, 1.
 " *rotundifolia*, VI, 12.
 " See also *microcentrum*.
Platamodes pennsylvanica, I, 1 and 3.
Platyphyllum. See *Cyrtophyllus*.
Podisma. See *Pezotettix*.
Pterolepis minutus, XI, 17.
Pterolepis. See *Anabrus* also.
Podura villosa, VI, 7.
Pterophylla. See also *Cyrtophyllus*, *Orchelimum*, *Xiphidium*.
Pyrgomorpha brevicornis, IV, 14.
Rhaphidophora. See *Ceuthophilus*, *Hadenæcus*.
Romalea microptera, III, 4.
Smythurus, VI, 7.
Spectrum. See *Diapheromera*, *Anisomorpha*.
Stauronotus, Elliotti, VIII, 11.
Stenobothrus admirabilis, IV, 13; V, 14.
 " *aequalis*, VI, 21.
 " *curtipennis*, VI, 15; VII, 10; X, 4; XII, 18.
 " *maculipennis*, X, 14; VI, 27.
 " *longipennis*, V, 15.
Stenopelmatus talpa, I, 10.
 " *fasciatus*, XIII, 7.
Stylopyga orientalis, I, 5, 6; VII, 12.
Tettigidea polymorpha, VII, 1.
Tettix lateralis, VIII, 7.
 " *ornata*, V, 1 and 2; XII, 19.
 " *Tetrix*, *Tettigidea*, *Batrachidea*.
Thamnotrizon dorsale, I, 11.
 " *scabralis*, XIII, 6.
 " *trilineatus*, VIII, 5.
Thrips (?) VI, 5, 9; XII, 11.
 " *cerealum*, XII, 9, 10.
Tomonotus. See *Edipoda tenebrosa*.
Tragocephala infusata, X, 8.
 " *viridifasciata*, V, 9.
Trichodectes equi, XII, 4.
 " *latus*, VI, 8.
 " *longicornis*, XII, 3.
Tridactylus terminalis, III, 6.
Udeopsylla robusta, VIII, 2.
Xiphidium brevipennis, IV, 11.
 " *fasciatum*, IV, 10.

ALPHABETICAL LIST OF SPECIES FIGURED.

- abbreviatus Gryllus, VII, 17.
 abbreviata Acheta. See Gryllus.
 abortiva Chlœaltis. See C. conspersa.
 acuminata Locusta. See Conocephalus, ensiger.
 aequalis CEdipoda, XIII, 8.
 " Locusta. See CEdipoda.
 " Gryllus. See "
 " Stenobothrus, VI, 21.
 agilis Pterophylla. See Orchelimum vulgare.
 alutaceum Acridium, VIII, 13; X, 13.
 americanum Acridium, I, 15.
 " Cyrtacanthacris. See Acridium.
 " Gryllus. See Acridium.
 americana Periplaneta, I, 2.
 " Blatta. See Periplaneta.
 americanus Gryllotalpa. See G. borealis.
 augustifolia Phaneroptera. See P. curvicauda.
 arenosa Tetrix. See T. Ornata.
 argulus Nirmus, XII, 1.
 atrox CEdipoda, VIII, 3.
 bipunctatus CEdipoda, IV, 5, 6.
 " Gryllus. See CEdipoda.
 bilineata Tetrix. See T. ornata.
 bivittatus Caloptenus, I, 16; V, 16.
 " Gryllus. See Caloptenus.
 bivittata Opomala, VI, 24-25; XI, 7.
 " Acridium. See Opomala.
 " Opsomala. See Opomala.
 bivittatum Spectrum. See Anisomorpha bu-
 prestoides.
 borealis Gryllotalpa, VII, 13,
 " Pezotettix, VI, 16.
 brevicornis Pyrgomorpha, IV, 14,
 " Opomala. See Pyrgomorpha.
 " Truxalis. See Pyrgomorpha.
 brevipennis Gryllotalpa. See G. borealis.
 " Xiphidium, IV, 11.
 brevipennis Dalmanina, VII, 14, 15.
 buprestoides Anisomorpha, I, 8.
 " Phasma. See Anisomorpha.
 carinata CEdipoda, X, 7.
 carlingiana " XII, 21.
 carolina " V, 3.
 " Locusta. See CEdipoda.
 caroliniana Locusta. "
 carolina, Mantis II, 1, 2, 3, &c.
 centurio Gryllus. See Romalea microptera.
 cerealeum Thrip, XII, 9, 10.
 cincta CEdipoda, XII, 13.
 coloradensis Phaneroptera, XI, 12.
 concavus Cyrtophyllus, IV, 15.
 concavum Platyphyllum. See Cyrtophyllus.
 concava Pterophylla. See Cyrtophyllus.
 conspersa Chlœaltis, VI, 11; X, 12.
 corallina Locusta. See CEdipoda, phoenicoptera.
 corallipes CEdipoda, III, 8. See also CEdipoda Halde-
 mannii.
 coriacea Phlœothrips, VI, 4.
 cristata Batrachidea, V, 8.
 " Tetrix. See Batrachidea.
 curtispennis Stenobothrus, VI, 15; VII, 17;
 X, 14; XII, 18.
 curtispennis Chlœaltis. See Stenobothrus.
 " Locusta. See Stenobothrus.
 curvicauda Phaneroptera, VII, 2, 3; V, 10.
 " Locusta. See Phaneroptera.
 cygni Docophorus, XII, 5.
 damnificum Acridium. See A. rubiginosum.
 differentialis Caloptenus, VIII, 12; IX, 4;
 XI, 6.
 discoidea CEdipoda, III, 7.
 dodgei Caloptenus, XI, 4, 5, 9.
 domesticus Gryllus, VI, 14.
 domestica Acheta. See Gryllus.
 dorsale Thamnotrizon, I, 17.
 dorsalis Tetrix. See Tettix ornata.
 elliotti Stauronotus, VIII, 11.
 ensiger Conocephalus, IV, 12.
 equi Trichodectes, XII, 4.
 eucrata CEdipoda, III, 2; V, 13; VI, 23.
 " Locusta. See CEdipoda.
 exiguus Nemobius, VII, 18.
 " Acheta. See Nemobius.
 falcicornis Philopterus, VI, 1.
 fasciatum Xiphidium, IV, 11.
 fasciatus Nemobius, III, 9, 10,
 " Gryllus. See Nemobius.
 " CEdipoda. See CEdipoda.
 " Stenopelmatus, XIII, 7.
 femorata Diapheromera, I, 7; X, 1.
 femoratus Bacunculus. See Diapheromera.
 femoratum Spectrum. See Diapheromera.
 femoratus Caloptenus. See C. bivittatus.
 femur-rubrum Caloptenus, V, 11; VIII, 2.
 " Acridium. See Caloptenus.
 flavicornis Acridium. See A. semi-rubrum.
 flavofasciatum Boopidou, VIII, 10.
 formosus Gryllus (?) IX, 5.
 frontalis Acridium, XI, 1.
 " Pezotettix, XIII, 2.
 fuliginosa Locusta, IX, 9.
 germanica Ectobia, I, 4.
 " Blatta. See Ectobia.
 gigantea Forficula, X, 2.
 gracile Orchelimum. See Xiphidium, fasciatum.
 gracilipes Ephippitytha, XI, 11.
 griseus Caloptenus, XII, 14.
 haldemanni Pterolepis. See Anabrus.
 haldemanni Anabrus, VII, 16.
 " CEdipoda, XIII, 3. }
 See also CEdipoda corallipes. }
 hirtipes Acrolophus, IX, 7.
 " Gryllus. See Acrolophus.
 hospes Acheta. See Nemobius fasciatus.

- infuscata Tragocephala, X, 8.
 " Gomphocerus. See Tragocephala.
 lapidicolus Ceuthophilus, VII, 4, 5.
 lapidicola Phalangopsis. See Ceuthophilus.
 " Raphidiphora. See Ceuthophilus.
 lateralis Tettix, V, 7.
 " Tettix. See Tettix.
 " Acridium. See Tettix.
 latipennis Locusta. See CEdipoda verruculata.
 latus Trichodectes, VI, 8.
 longicornis Trichodectes, XII, 3.
 longipennis Gryllotalpa, I, 12.
 " Stenobothrus, V, 15.
 luctuosus Gryllus, IX, 10.
 " Acheta. See Gryllus.
 maculatus Ceuthophilus, III, 5.
 maculata Ephippigera. See Ceuthophilus.
 " Phalangopsis. See Ceuthophilus.
 " Raphidiphora. See Ceuthophilus.
 maculipennis Stenobothrus, VI, 27; X, 14.
 magnus, Brachypeplus, VII, 14.
 maritima CEdipoda, XII, 17.
 " Locusta. See CEdipoda.
 marmorata CEdipoda, VII, 2.
 " Locusta. See CEdipoda.
 microptera Romalea, III, 4.
 minor Labia, X, 3.
 " Forficula. See Labia.
 minuta Labia, I, 10.
 minutus Pterolepis, XI, 17.
 minutus Anabrus. See also Pterolepis.
 missouriensis Mantis, XIII, 11.
 montana CEdipoda, XII, 12.
 mucronata Copiophora, VII, 8; VIII, 14.
 " Copiophora. See Copiophora.
 nebulosa Locusta. See CEdipoda sordida.
 nebrascensis Pezotettix, XII, 2.
 neglectus Gryllus, X, 8.
 nietanus Tomonotus. See CEdipoda tenebrosa.
 nigrum Boopidon. See B. nubilum.
 niveus CEdipoda, IV, 1, 2.
 " Gryllus. See CEdipoda.
 nubilum Boopidon, XIII, 10.
 nubilus Gryllus. See Boopidon.
 obesa Pezotettix, XI, 13, 14.
 oblongifolia Phylloptera, IV, 4; X, 5; and
 XIII, 1.
 oblongifolia Locusta. See Phylloptera.
 obscurum Acridium, V, 2.
 obscurus Gryllus. See Acridium.
 obscurus Oxycoryphus, XIII, 13.
 occidentalis Locusta, XI, 15.
 orientalis Stylopyga, I, 5, 6; VII, 12.
 " Blatta. See Stylopyga.
 ornata Tettix, V, 1, 2; XII, 19.
 ornatum Acridium. See Tettix.
 " Tettix. See Tettix.
 pallidum Menapon, XII, 2.
 pallidipalpis Decticus, IX, 8.
 parvipennis Tettix. See Tettigidea, Polymorpha.
 pavonius Pediculus. See Phyllopterus falcicornis.
 pellucida CEdipoda, XII, 20.
 pennsylvanica Platamodes, I, 1, 3.
 " Blatta. See Platamodes.
 pennsylvanicus. See Gryllus.
 phœnicoptera CEdipoda, V, 4.
 picta Pezotettix, VIII, 4.
 piliferus, Haematopinus, XII, 6.
 polymorpha Tettigidea, VII, 1.
 " Tettix. See Tettigidea.
 pruinosa CEdipoda. See CEdipoda trifasciata.
 pulchellus Phyllopalpus, VI, 22.
 punctulatus CEdipoda. See CEdipoda, bi-
 punctatus.
 punctulatus Cryptocercus, VI, 20.
 quadrimaculata Tettix. See T. Ornata.
 retinervis Microcentrum, IV, 3.
 " Phylloptera. See Microcentrum.
 robusta Udeopsylla, VIII, 2.
 " Daihinia. See Udeopsylla.
 " Phalangopsis. See Udeopsylla.
 rotundifolia Phylloptera, VI, 12.
 rugosa CEdipoda, XII, 8.
 rubiginosum Acridium, V, 5.
 rusticum (Burm. only) Acridium. See A. Alu-
 taceum.
 radiata Tragocephala. See T. viridifasciata.
 saccharina Lepisma, VI, 6.
 saltator Orocharis, III, 11, 12.
 sayii Bacteria. See Diapheromera femorata.
 scabricollis Thamnotrizon, XIII, 6.
 scudderi Camptonotus, VIII, 15.
 semi-rubrum Acridium, XIII, 14.
 servilis Acheta. See Nemobius vittatus.
 simplex Anabrus, IX, 1.
 sordida CEdipoda, X, 11.
 " Tettix. See T. ornata.
 spretus Caloptenus, VIII, 1; XIII, 15.
 subterraneus Hadenæus, VIII, 6.
 " Raphidiphora. See Hadenæus.
 sulphurea CEdipoda, V, 6.
 " Locusta. See CEdipoda.
 " Gryllus. See CEdipoda.
 suis Haematopinus, XII, 4.
 talpa Stenopelmatus, I, 10.
 tenebrosa CEdipoda, IX, 2.
 terminalis Tridactylus, III, 6.
 torvum Acridium. See C. Alutaceum.
 trifasciata CEdipoda, IX, 6.
 " Gryllus. See CEdipoda.
 trilineatus Thamnotrizon, VIII, 5.
 " Decticus. See Thamnotrizon.
 tschivavensis Ephippigera, XI, 15.
 tuberculatum Acridium. See A. disoidea.
 turbullii Caloptenus, XI, 10.
 uhleri Ceuthophilus, VIII, 8.
 undulata CEdipoda, XII, 15.
 unicolor Pezotettix, XIII, 4.
 variabilis Machilis, VI, 2, 3.
 verruculata CEdipoda, X, 6.
 " Locusta. See CEdipoda.
 " Phrynotettix, VI, 25.
 villosa Podura, VI, 7.
 virescens Brachypeplus, IX, 3.
 virginianum Acridium. See Tragocephala viri-
 difasciata.
 virginiana CEdipoda. See viridifasciata.
 viridis Caloptenus, XI, 3.
 viridis Chloealetis, X, 5.
 " Stenobothrus. See Chloealetis.
 viridifasciata Tragocephala, V, 9.
 vittatus Nemobius, III, 9, 10.
 vulgare Orchelimum, IV, 7, 8; VII, 6.
 wyomingensis Mesops, IV, 9; XI, 8.
 " Opomala. See Mesops.



LIST OF DESIDERATA TO FINISH THE WORK.

Hapithus agitator, Uhler, (Md.)	Tettix rugosa, Scudd., (Mass.)
Gryllus angustus, Scudd., (Mass.)	Batrachidea carinata, Scudd., (Cal.)
Xya apicalis, Uhler, (South Western States.)	Thespis, (Mantis,) parva, Drury, pl. II, 39, (Amer.)
Tridactylus minutus, Scudd., (Ill.)	Phibalosoma ploiaria, Western, (Western State.)
Cyphoderris monstrosa, Uhler, (Oregon.)	Temnopteryx aeropeltiformis, Burm., (U. S.)
Platyphyllum perspicillatum, Serv. and Uhler in Harris (U. S.)	" virginiana, Burm., (Va.)
Thyreonotus dorsalis, Scudd., (Mass. and Md.)	Ischnoptera Uhleriana, Sauss., (U. S. Pa.) or any in U. S.
Orchesticus americanus, Sauss., (Tenn.)	Platamodes unicolor, Scudd., (Mass.)
Gryllacris carolinensis, Gerst., (Car.)	Ectobia lithophila, Scudd. (Mass.)
Tropidischia xanthostoma, Scudd., (Cal.)	Paratrapes histrio, Saus., (U. S.)
Polisma (Acridium,) borkii Stål, (Cal.)	Pycnoscelis obscurus, Scudd., (Mass.)
Acridium flavofasciatum, Thomas, (Col.)	Brachylabris maratima, Dohrn. (Southern State.)
Arcyptera lineata, Scudd. (Mass.)	Psolidophora brunneipennis, Serv., (Pa. and Va.)
" platyptera, Scudd., (New Eng.)	" bipunctata Dohrn, (Mass.)
" gracilis, (Red River, Brit. Am.)	Forficula pulchella (Serv.) (Magura.)
Tragocephala radiata, Erichs, (Mass.) (prob. syn. infusata. ?.)	Thrips (Limothrips) tritici, from Pack.
Tettix cucullata, Burm. (Mass.)	Thrips (Heliothrips) hæmorrhoidalis, Burm. See Pack.
" granulata, Scudd., "	
" triangularis, " "	

Authenticated specimens of any of the above-mentioned Orthoptera will be most thankfully received, and, after being figured, will be returned if required. Also, any other new species not mentioned in the list of insects figured, if sent to the Author, will be figured and credited to the donor or sender and placed in an extra plate of addenda, whenever a sufficient number of specimens are received to fill a plate.

ERRATA:

Page V, and Plates III and IV; for "*Æ*" in *Æcanthus* and *Ædipoda*, read *Æ*.
 Plate IV, 10; for "*Xyphidium*," read *Xiphidium*.
 Plate VII, 1; for "*parvepennis*," read *parvipennis*.
 Page 2, line 16; for "*larvae*," read *larvæ*.
 Page 2, line 33; insert commas after "*food*" and "*Indians*."
 Page 2, line 35; dele "*which*" after U. S. A.
 Page 3, last line; for "*Chalcididea*," read *Chalcididae*.
 Page 6, after swan; for "*Docophorus*," read *Docophorus*.
 Page 9; for "*angustifolia*," read *angustifolia*.

